

# Quality of care assessment for people with multimorbidity

J. M. Valderas<sup>1,2,3,4</sup> , J. Gangannagaripalli<sup>1</sup>, E. Nolte<sup>5</sup>, C. M. Boyd<sup>6,7</sup>, M. Roland<sup>8</sup>, A. Sarria-Santamera<sup>9,10,11,12</sup>, E. Jones<sup>1</sup> & M. Rijken<sup>13,14</sup>

<sup>1</sup>Health Services & Policy Research, Exeter Collaboration for Academic Primary Care (APEX), University of Exeter, Exeter, UK; <sup>2</sup>Institut Universitari d'Investigació en Atenció Primària (IDIAP Jordi Gol), Barcelona, Spain; <sup>3</sup>Threads & Yarns International Multimorbidity Research Network, Exeter, UK; <sup>4</sup>National Institute for Health Research (NIHR) Collaboration for Leadership in Applied Health Research and Care (CLAHRC) South West, University of Exeter, Exeter; <sup>5</sup>London School of Hygiene & Tropical Medicine, London, UK; <sup>6</sup>The Johns Hopkins School of Medicine; <sup>7</sup>The Johns Hopkins Center on Aging and Health, Baltimore, MD, USA; <sup>8</sup>Department of Public Health and Primary Care, University of Cambridge, Cambridge, UK; <sup>9</sup>Escuela Nacional de Sanidad, Instituto de Salud Carlos III, Madrid; <sup>10</sup>Universidad de Alcalá, Alcalá de Henares, Madrid; <sup>11</sup>Instituto Mixto de Investigación IMIENS, Escuela Nacional de Salud, Universidad Nacional de Educación a Distancia; <sup>12</sup>REDISSEC, Spanish Network of Health Services and Chronic Patients, Madrid, Spain; <sup>13</sup>Netherlands Institute for Health Services Research (NIVEL), Utrecht, The Netherlands; and <sup>14</sup>Department of Health and Social Management, University of Eastern Finland, Kuopio, Finland

**Content List** – Read more articles from the symposium: “Multimorbidity research at the cross-roads: developing the evidence for clinical practice and health policy”.

**Abstract.** Valderas JM, Gangannagaripalli J, Nolte E, Boyd C, Roland M, Sarria-Santamera A, Jones E, Rijken M (University of Exeter, Exeter, UK; Institut Universitari d'Investigació en Atenció Primària (IDIAP Jordi Gol), Barcelona, Spain; Threads & Yarns International Multimorbidity Research Network, Exeter, UK; London School of Hygiene & Tropical Medicine, London, UK; The Johns Hopkins School of Medicine; The Johns Hopkins Center on Aging and Health, Baltimore, MD, USA; University of Cambridge, Cambridge, UK; Instituto de Salud Carlos III, Madrid, Spain; Universidad de Alcalá, Alcalá de Henares; Universidad Nacional de Educación a Distancia, Madrid, Spain; Netherlands Institute for Health Services Research (NIVEL), Utrecht, The Netherlands; University of Eastern Finland, Kuopio, Finland). Quality of care assessment for people with multimorbidity. *J Intern Med* 2019; **285**: 289–300.

Multimorbidity, the simultaneous presence of multiple health conditions in an individual, is an increasingly common phenomenon globally. The systematic assessment of the quality of care delivered to people with multimorbidity will be key to informing the organization of services for meeting their complex needs. Yet, current assessments tend to focus on single conditions and do not capture the complex processes that are required for providing care for people with multimorbidity. We conducted a scoping review on quality of care and multimorbidity in selected databases in June 2018 and identified 87 documents as eligible for review, predominantly original research and

reviews from North America, Europe and Australasia and mostly frequently related to primary care settings. We synthesized data qualitatively in terms of perceived challenges, evidence and proposed metrics. Findings reveal that the association between quality of care and multimorbidity is complex and depends on the conditions involved (quality appears to be higher for those with concordant conditions, and lower in the presence of discordant conditions) and the approach used for measuring quality (quality appears to be higher in people with multimorbidity when measured using condition/drug-specific process or intermediate outcome indicators, and worse when using patient-centred reports of experiences of care). People with discordant multimorbidity may be disadvantaged by current approaches to quality assessment, particularly when they are linked to financial incentives. A better understanding of models of care that best meet the needs of this group is needed for developing appropriate quality assessment frameworks. Capturing patient preferences and values and incorporate patients' voices in the form of patient-reported experiences and outcomes of care will be critical towards the achievement of high-performing health systems that are responsive to the needs of people with multimorbidity.

**Keywords:** comorbidity, health systems performance assessment, multimorbidity, patient reported outcomes measures, patient safety, quality of health care.

## Introduction

Chronic conditions contribute to a large proportion of the morbidity burden and pose a major challenge to health systems worldwide [1]. Response to chronic conditions is frequently complicated by multimorbidity, the simultaneous presence of multiple health conditions in an individual [2–5]. Multimorbidity challenges usual care delivery, which is frequently structured around pathways of care for single diseases [6–10]. Key principles have been proposed for the design of high-performing health systems that meet the complex needs of people with multimorbidity, ranging from patient and caregiver engagement, to information systems, alignment of funding and incentives [11, 12]. Sustainable models of integrated care for multimorbidity currently being explored [13]. However, the evidence for how to effectively improve health outcomes for people with multimorbidity remains patchy [10, 14, 15], as confirmed by an updated systematic review [16]. A recent randomized evaluation of a complex multidimensional intervention simultaneously targeting medicines management, mental health and patient centredness has further highlighted the continued challenge of demonstrating evidence of effect in this complex population [17].

Efforts to improve the outcomes of care for people with multimorbidity can be supported by the rigorous monitoring and evaluation of service delivery as part of a health system performance framework to inform evidence-based decision-making [18–21]. There has been growing interest in the systematic evaluation of the quality of health care (the degree to which health services for individuals and populations increase the likelihood of desired outcomes and are consistent with current professional knowledge) [19, 22–25]. This has included considerable work into the development and use of quality indicators for a range of prevalent conditions, such as ischaemic disease, stroke, COPD, diabetes and cancer, with some countries such as the United Kingdom or the United States linking performance based on these indicators to financial and nonfinancial incentives in an effort to improve the quality of care [19, 26, 27].

It has become increasingly clear, however, that a continued focus on the quality of care for single conditions fails to capture the complex processes required for providing care across conditions, nor does it provide the right stimulus to improve those service delivery components that are core to

providing high-quality care for people with multimorbidity, such as coordination and integration of care [6, 9, 28].

Overall, there remains a need to systematically bring together the existing evidence base on efforts to assess the quality of care delivered to people with multimorbidity to help inform the development of an assessment framework that can then inform decision-making on the organization and delivery of care that better meets the complex needs of people with multimorbidity. This paper seeks to contribute to this process by means of a scoping review that (i) explores how this issue has been framed in the literature, (ii) examines the empirical evidence of the association between quality of care and multimorbidity and (iii) assesses metrics and frameworks that have been proposed for the evaluation of the quality of care delivered to people with multimorbidity.

## Methods

We conducted a scoping review of the literature on multimorbidity and healthcare performance assessment focussing on quality of healthcare processes and outcomes. We selected this approach as an established method for clarifying conceptual boundaries and mapping out research areas that have not yet been extensively reviewed, and that are of complex and heterogeneous nature [29, 30].

We searched the following databases: OVID including MEDLINE, EMBASE and Health Management Information Consortium (which includes the English Department of Health's Library and Information Services (DH-Data) and the King's Fund Information and Library Service), PubMed and the bibliographic database on multimorbidity maintained at the Health Services & Policy Research Group at the University of Exeter, which is updated weekly from ISI Web of Science and Google Scholar alerts for documents using the term 'multimorbidity'. We developed bespoke search strategies for each database using Boolean operators to link two main blocks: multimorbidity and healthcare performance. We used the overarching term of 'healthcare performance' rather than the more narrow notion of 'quality of care processes and outcomes' to ensure the searches capture the wide range of work that may be of relevance to this study. This is based on our previous experience of conducting reviews of quality of care indicators that found that terms 'quality' and 'performance' are often used

interchangeably, although the latter is typically understood as a broader, multidimensional concept that, in addition to quality, also includes dimensions of equity and efficiency [31]. While we recognize these important conceptual differences, in this paper, we will use the terms interchangeably also, reflecting the varying ways authors of papers included in this review have used these terms.

The search was implemented on 15 June 2018. We did not impose any restrictions on publication date, journal and type of publication or language. All citations were imported into the bibliographic manager EndNote. Duplicate citations were first removed automatically and subsequently through a manual process when needed.

A three-stage screening process was used to assess the relevance of studies identified in the search. Studies were eligible for inclusion if they made any reference to the assessment of healthcare quality for people with multimorbidity, with a specific focus on processes and outcomes of care. For the first level of screening, only the titles of citations were reviewed with a sensitive approach in which only documents whose scope was clearly outside the scope of this review were excluded. Title screening was piloted by three authors (JMV, JG and EJ) with 50 randomly selected titles in order to ensure consistent application of the eligibility criterion and then was subsequently applied independently by two reviewers (JG and EJ). In cases of disagreement, the document was included in the next stage. The second level involved abstract review of documents deemed potentially eligible in the previous step using the same inclusive and sensitive approach. The process was replicated for abstracts (pilot with 20 abstracts). In the third step, full texts of the documents deemed potentially eligible were screened (pilot with 5 papers). Disagreement was resolved at this stage by consensus. The characteristics of each full-text article were extracted by two reviewers (JG and EJ) using a standardized template. Based on a predefined framework, a narrative synthesis of the information contained in the included documents was conducted initially by two authors (JG and JMV) for comment and review by all authors. The proposed framework included problem framing (justification of a focus on multimorbidity in the evaluation of healthcare quality); evidence (empirical data for the association between multimorbidity and the quality of process and outcomes of care); and measurement (metrics and frameworks that have been proposed

for the evaluation of performance in the presence of multimorbidity). Formal assessment of the quality of included studies was deemed inappropriate given the scope of the review and the broad range of types of articles retrieved.

## Results

### *Search results*


The search retrieved 435 documents after removal of duplicates (Fig. 1), and after eligibility screening, a total of 87 documents were finally included [6–9, 11, 13, 16, 28, 32–110] (Appendix S1).

The literature reviewed included a wide range of documents, including original studies using qualitative and quantitative research methods, systematic reviews, policy briefs, editorials and commentaries, reports, and other (Table 1). The majority of documents originated in the United States, Canada, selected European countries (UK, Netherlands and Ireland), New Zealand and Australia and were published in the last 5 years (Appendix).

### *Framing of the problem and perceived challenges*

The literature reviewed justifies the need to focus on the evaluation of quality of care delivered to people with multimorbidity on grounds of the large numbers of those affected, and the impact of multimorbidity on healthcare processes and outcomes [103]. Concerns about the rising prevalence of multimorbidity are largely attributed to an increased prevalence of individual chronic conditions and to the association of multimorbidity with increasing age [37].

People with multimorbidity face a higher risk of complications of medical care, including pharmacological interactions and adverse drug events, avoidable admissions, and misalignment of multiple care plans proposed by different health professionals. These are perceived to be the attributable to higher service utilization in this population group (both more frequent and more varied utilization across multiple settings, and polypharmacy) as well as the intrinsic complexity of their clinical management [37, 39, 44, 66, 80]. High levels of service utilization are generally seen as the key determinant of increased healthcare costs, poor patient satisfaction and, potentially, also a contributor to adverse health outcomes, which include poor quality of life, reduced ability to work and



**Fig. 1** PRISMA Flowchart of the study selection process.

employability, and increased disability and mortality [84, 86, 90].

There is consensus in the reviewed literature that the main challenge posed by multimorbidity for achieving high healthcare performance is the current organization of health care following a 'disease-oriented' model. This has broad implications, ranging from care financing and reimbursement to the degree of applicability of current clinical practice guidelines to this patient group [89]. Disease orientated care results in fragmentation and lack of coordination and continuity of care, making people with multimorbidity particularly vulnerable during transitions of care [63]. The literature supports the key role played by primary care's patient focussed approach in contributing to both coordination and continuity of care [33, 51]. Lack of robust evidence on the most appropriate care for people with different

multimorbidity profiles is recognized as a challenge for the provision of efficient and effective care [43]. The usually limited involvement of individuals in decision-making is perceived as a significant challenge for people with multimorbidity, as continued uncertainty about best management approaches makes effective patient engagement crucial [8].

#### *The association of multimorbidity and quality of care: empirical evidence*

Ricci-Cabello and colleagues have highlighted the complex association between quality of care and multimorbidity in their recent review, which found that the direction of the association seemed to depend on the constructs used for multimorbidity and quality assessment and their operationalization [88]. The quality of care appeared to be higher when quality was measured using condition/drug-

**Table 1** Characteristics of included documents (n = 87)

| Characteristic       | n (%)   |
|----------------------|---------|
| Year                 |         |
| 2006–2010            | 14 (16) |
| 2011–2015            | 38 (44) |
| 2016–2018            | 35 (40) |
| Type of document     |         |
| Original research    | 38 (44) |
| Review               | 22 (25) |
| Policy brief         | 3 (3)   |
| Other                | 24 (28) |
| Setting <sup>a</sup> |         |
| Primary Care         | 44 (47) |
| Other setting        | 20 (21) |
| Nonspecific          | 30 (32) |
| Country <sup>a</sup> |         |
| USA                  | 38 (31) |
| UK                   | 15 (12) |
| Canada               | 9 (7)   |
| Australia            | 8 (7)   |
| Germany              | 7 (6)   |
| Other                | 46 (37) |

<sup>a</sup>Categories exceed 100% as categories are not mutually exclusive. See Appendix for full details of included studies.

specific process or intermediate outcome indicators, and worse when quality was measured using patient-centred reports of experiences of care [88]. Of note, studies that explored the related construct of comorbidity (which considers the presence of conditions in relation to an index disease) found that care quality may be higher for those with concordant conditions (e.g. those sharing a common pathophysiological pathway and therefore more likely to benefit from the same clinical management, such as hypertension, ischaemic heart disease and diabetes), and lower in the presence of discordant conditions (those not sharing a common pathophysiological pathway, such as COPD and diabetes) [88, 110].

Panagioti *et al.* focussed specifically on safety in people with multimorbidity, finding that patient safety events (and their type) varied by the nature of multimorbidity [85]. Thus, people with physical and mental health conditions were found to be at a higher risk of safety incidents than those with multimorbidity that did not involve mental health. Multimorbidity

was also associated with increased risk of incidents that resulted in adverse outcomes [85].

#### *Quality metrics and assessment frameworks for care for people with multimorbidity*

Approaches to the evaluation of quality of care for people with multimorbidity in the reviewed literature frequently relies on aggregating disease-specific indicators for the quality of processes and outcomes of care [62], which are typically derived from single-disease-oriented guidelines [6]. This additive model that considers quality of care for multimorbidity as the sum of estimates of quality of care for each individual condition is viewed critically [44], given the lack of robust empirical evidence supporting the validity of this approach [7]. Disease-oriented guidelines may have limited applicability to people with multimorbidity [90], given their reliance on clinical trials which typically exclude medically complex patients or people undergoing multiple medical interventions. However, such patients are most commonly seen in clinical practice [89]. The additive approach does not account either for the potential of interactions between different treatments, between treatments and diseases (with the first complicating the prognosis and management of the latter) and between diseases, with potentially harmful consequences [68]. The additive approach also means that quality of care for some diseases may be given priority when there is wide variation in the number of indicators available for each condition [91].

The reviewed literature supports the need for the development of performance measures that are specific for multimorbidity [53, 84] or nonspecific but robust in the presence of multimorbidity [7, 9]; rely on data from the electronic health record [39]; and include outcomes and processes of care, where there is evidence that the latter lead to improved outcomes [56]. The literature identifies a number of domains, and related measures, that broadly focus on areas reflecting the deficiencies in the provision of health care for people with multimorbidity that we have described above, and the outcomes of interventions targeting multimorbidity [16] (Box 1). However, much of the literature focuses on individual domains rather than bringing them together as part of a comprehensive assessment framework.

Experience in the development of multimorbidity-specific performance measures is still limited [87].

**Box 1.** Domains relevant to quality of care and performance assessment in people with multimorbidity

|   |
|---|
| Process of care   |
| Continuity  |
| Coordination  |
| Comprehensiveness   |
| Patient centredness   |
| Preferences elicitation   |
| Prioritization  |
| Individualized goal setting   |
| Self-efficacy   |
| Management of life style factors  |
| Management of specific diseases   |
| Medicines management  |
| Use of health services  |
| Experience of care and satisfaction   |
| Experiences of care   |
| Satisfaction with care  |
| Outcomes of care  |
| Patient-reported outcomes (symptoms, functioning, health-related quality of life) |
| Adverse events  |

The validity of such measures is contingent on the evidence supporting them and there remains paucity of research on best clinical approaches for people with multimorbidity [74]. However, this is changing rapidly as an increasing body of research is being developed to address this gap [16].

A number of initiatives for the development of comprehensive frameworks for performance assessment for people with multimorbidity are identified in the literature. The Organisation for Economic Co-operation and Development (OECD) is developing survey-based patient-reported indicators for capturing the experience and outcomes of care for patients with one or more chronic conditions [82]. Two core principles for the development of these indicators are patient involvement and the enablement of providers to use information for quality improvement and

shared decision-making. In parallel, the International Consortium for Health Outcomes Measurement, an independent consortium which the explicit goal of improving health system performance through standardized measurement, reporting and use of patient outcomes, is developing a core set of outcomes for overall adult health with the explicit goal of ensuring relevance to people with multimorbidity [111, 112]. Although these two initiatives were developed independently, they are increasingly being aligned to avoid duplications of efforts [112].

At national level, the Department of Health and Human Services (DHHS) of the US Federal Government has acknowledged that the promotion of best practices in caring for individuals with multimorbidity requires specific performance measures that consider the complex and dynamic nature of care for these patients [86]. A measurement framework to facilitate the development and refinement of such measures has been proposed in collaboration with the National Quality Forum (NQF). The framework is centred around patient and family goals and preferences for care in the context of multiple care sites and providers, the type of care they are receiving and considers the following priority domains for healthcare quality measurement, including (i) optimizing function, maintaining function or preventing further decline in function; (ii) seamless transitions between multiple providers and sites of care; (iii) patient important outcomes (includes patient-reported outcomes and relevant disease-specific outcomes); (iv) avoiding inappropriate, nonbeneficial care, including at the end of life; (v) access to a usual source of care, transparency of cost (total cost); (vi) shared accountability across patients, families and providers; and (vii) shared decision-making [53, 56].

## Discussion

This review has identified a number of documented efforts to advance thinking, evidence and methods in the area of quality of care for people with multimorbidity. This emerging body of evidence and methods can be further developed towards a comprehensive assessment framework for an effective health system response to the rising burden of multimorbidity.

We used a scoping review to capture the complex and heterogenous body of evidence around multimorbidity and healthcare quality. We sought to be

inclusive in the type and nature of documents considered for review using very broad search terms. Clearly, any such approach may still miss relevant literature. More importantly perhaps, we will have not captured ongoing work on care quality and models for people with multimorbidity, which remains an emergent field, in particular ongoing work on indicator development. We recognize this limitation arguing that it would have required a different approach to the review and which was not feasible within the scope of this study. We believe, however, and within these limitations, that the retrieved literature gives a broad perspective of the current state of the art of advances in this area.

Our review has identified a number of important lessons around the systematic assessment of the quality of processes and outcomes of care for people with multimorbidity.

First, although there is evidence that multimorbidity may be associated with higher performance as measured by disease-specific indicators, current approaches to performance assessment may

disadvantage people with multimorbidity, particularly for patients with discordant conditions. Available condition-specific indicators do not provide the right incentives for managing patients with multimorbidity and may act as a barrier for providing best care. Adjusting quality of care for multimorbidity (risk adjustment) or even incentivizing the delivery of care for people with multimorbidity offers only partial solutions as they would not need to address the core problem of the validity of the measures in this group of patients. Appropriate quality measures for multimorbidity are needed, and the frameworks reviewed in this paper are steps in this direction, while still very much in need for further development and support by evidence. Research on the burden of discordant conditions is needed for targeting those patients that may benefit most from this expanded approach.

Second, measures of quality of care need to be consistent with the models of care, their processes and their relevant outcomes. Epidemiological transitions across the globe made it necessary to adapt models of care essentially oriented to an acute




Fig. 2 Models of care as informed by models of disease.

disease model (linear approach focussing on a single aetiological agent and the delivery of a single treatment) to effectively respond to chronic conditions (iterative approach dealing with multiple aetiological agents and multiple management options). A similar transition is needed from a single-disease model to a multimorbidity model. Such a model (and the assessment of its performance) has to account for the need to integrate care across conditions and providers and recognize the importance of patient-centred care with explicit goal setting and prioritization [7, 12, 92, 109, 113–115] (Fig. 2).

Third, the assessment of the quality of primary care should be at the core of evaluations of the care that people with multimorbidity receive. Transitions between providers and between episodes of care are critical to the needs of people with multimorbidity, requiring systematic coordination, continuity and comprehensiveness. Together with first contact care and person focus, these are also core functions of primary care [22, 117]. This well-established person focussed approach to health-care delivery can be considered the core model of care on which to base further developments oriented to improving care for people with multimorbidity [12, 22, 117], as the primary care focus of both the OECD PaRIS and ICHOM initiatives demonstrate.

Fourth, person-centred care should be a guiding principle for the development of assessment frameworks. People centredness, a core value of health systems, acknowledges that individual service users should be the key stakeholders [118, 119]. Their values, goals and priorities should shape care delivery and individual care plans, and this should be reflected accordingly in quality indicators. It has been proposed that making care more person centred may also counter the care fragmentation, which is particularly detrimental to care of patients with multimorbidity, while increasing patient satisfaction [90].

Considering the evidence reviewed here, we identify two priority areas for further research and development. First, there is an urgent need to establish how to enable the routine collection of patient evaluations of health and health care using patient-reported experience and outcome measures (PREMS and PROMs) and to incorporate these into comprehensive assessment frameworks [21, 106, 120–124]. Second, there is a need to

advance approaches for the measurement of the role of service users (and their carers) as active partners in service delivery. This is notoriously difficult to capture in current information systems and developing the methods for best documenting and evaluating performance on core aspects such as explicit goal setting and prioritization should be a research priority [116, 125].

### Conclusion

Single-disease approaches to the measurement of quality of care for people with multimorbidity do not capture the complexity of the processes involved in meeting the complex needs of this population. This scoping review has identified important avenues for the further development of approaches for the systematic assessment of the quality of care for people with multimorbidity, but also highlighted the need for a critical shift in our understanding of the underlying models of care that can best meet the needs of this group for developing the evidence base. Assessment frameworks that capture patient preferences and values and incorporate patients' voices in the form of patient-reported experiences and outcomes of care will be critical for making progress towards the achievement of high-performing health systems.

### Authors' contributions

JMV is the grantor. JMV designed the concept of the paper, which was agreed with CB, EN, MaR, MiR and ASS. EJ, JG and JMV had full access to all of the data in the study, implemented the search strategy, applied eligibility criteria and extracted the information. JG and JMV conducted the analysis. ASS, CB, EN, JMV, MaR and MiR contributed to the Session 'Multimorbidity and Health Policy' of the International Symposium 'Multimorbidity research at the cross-roads: developing the evidence for clinical practice and health policy' that took place on 21/05/2018 at the Nobel Forum, Karolinska Institutet, Stockholm, Sweden co-organized by Journal of Internal Medicine, Karolinska Institutet, University of Exeter, Threads & Yarns Collaboration, and KI Strategic Research Area in Epidemiology (SfoEpi), and with Laura Fratiglioni and JMV as scientific co-chairs (for programme, presentations and lessons learned, see <http://www.multimorbidity2018-stockholm.se> and <https://wol-prod-cdn.literatumonline.com/pb-assets/assets/13652796/Conference%20report%20Multimorbidity.pdf>) and to a subsequent

workshop where core aspects relevant to the development of the study were discussed. JMV drafted the first version of the manuscript, and all the authors (ASS, CB, EN, EJ, JG, JMV, MaR and MiR) revised subsequent drafts critically for important intellectual content and approved the final draft for publication.

### Funding

Journal of Internal Medicine, Karolinska Institutet Strategic Research Area in Epidemiology (SfoEpi).

### Conflict of interest statement

JMV has contributed as an advisor to the development of the OECD PARIS initiative. JMV is the chair of the ICHOM panel for Overall Adult Health core set.

### References

- WHO. *World Health Statistics 2018: Monitoring Health for the SDGs, Sustainable Development Goals*. Geneva: World Health Organization, 2018.
- Valderas JM, Starfield B, Sibbald B, Salisbury C, Roland M. Defining comorbidity: implications for understanding health and health services. *Ann Fam Med* 2009; **7**: 357–63.
- Violán C, Foguet-Boreu Q, Roso-Llorach A *et al.* Burden of multimorbidity, socioeconomic status and use of health services across stages of life in urban areas: a cross-sectional study. *BMC Public Health* 2014; **14**: 530.
- Calderón-Larrañaga A, Vetrano DL, Ferrucci L *et al.* Multimorbidity and functional impairment: bidirectional interplay, synergistic effects and common pathways. *J Intern Med* 2019; **285**: 255–71.
- Mounce LTA, Campbell JL, Henley WE, Tejerina Arreal MC, Porter I, Valderas JM. Predicting incident multimorbidity. *Ann Fam Med* 2018; **16**: 322–9.
- Boyd CM, Darer J, Boulton C, Fried LP, Boulton L, Wu AW. Clinical practice guidelines and quality of care for older patients with multiple comorbid diseases: implications for pay for performance. *JAMA* 2005; **294**: 716–24.
- Tinetti ME, Fried TR, Boyd CM. Designing health care for the most common chronic condition—multimorbidity. *JAMA* 2012; **307**: 2493–4.
- Boyd CM, Fortin M. Future of multimorbidity research: how should understanding of multimorbidity inform health system design? *Public Health Rev* 2010; **32**: 451.
- Valderas JM. Multimorbidity, not a health condition or complexity by another name. *Eur J Gen Pract* 2015; **21**: 213–4.
- Muth C, Blom JW, Smith SM *et al.* Evidence supporting the best clinical management of patients with multimorbidity and polypharmacy: a systematic guideline review and expert consensus. *J Intern Med* 2019; **285**: 272–88.
- Mossialos EAO, Roland R, Abrams M *et al.* Designing a high-performing health care system for patients with complex needs. Ten recommendations for policymakers. 2017, The Commonwealth Fund International Experts Working Group on Patients with Complex Needs.
- Mai S, Adam S, Ruth T, Rebecca F, Catherine T, Sarah D. *Briefing: Understanding the Health Care Needs of People with Multiple Health Conditions*. London: The Health Foundation, 2018.
- Leijten FRM, Struckmann V, van Ginneken E *et al.* The SELFIE framework for integrated care for multi-morbidity: development and description. *Health Policy* 2018; **122**: 12–22.
- WHO. High-Level Regional Meeting Health Systems Respond to NCDs: Experience in the European Region Sitges, Spain, 16–18 April 2018 Outcome statement. 2018, World Health Organization.
- Boyd CM, Kent DM. Evidence-based medicine and the hard problem of multimorbidity. *J Gen Intern Med* 2014; **29**: 552–3.
- Smith SM, Wallace E, O'Dowd T, Fortin M. Interventions for improving outcomes in patients with multimorbidity in primary care and community settings. *Cochrane Database Syst Rev* 2016; **3**: Cd006560.
- Salisbury C, Man MS, Bower P *et al.* Management of multimorbidity using a patient-centred care model: a pragmatic cluster-randomised trial of the 3D approach. *Lancet* 2018; **392**: 41–50.
- WHO. *Health System Performance Assessment: A Tool for Health Governance in the 21st Century*. Copenhagen, Denmark: World Health Organization, 2012.
- Smith PC, Mossialos E, Leatherman S, Papanicolas I (eds). *Performance Measurement for Health System Improvement: Experiences, Challenges and Prospects*. Cambridge: Cambridge University Press, 2009.
- Smith P. *Health System Performance Assessment*. Luxembourg: Publications Office of the European Union, 2014.
- Greenhalgh J, Dalkin S, Gibbons E *et al.* How do aggregated patient-reported outcome measures data stimulate health care improvement? A realist synthesis. *J Health Serv Res Policy* 2018; **23**: 57–65.
- WHO. *A Vision for Primary Health Care in the 21st Century - Towards Universal Health Coverage and the Sustainable Development Goals in Technical Series on Primary Health Care*. Geneva: World Health Organization, 2018.
- WHO. Primary health care: transforming vision into action. Operational Framework. Draft for consultation., in *Technical Series on Primary Health Care*. Geneva: World Health Organization, 2018.
- Academy A. A model for measuring quality care, in Online library of Quality, Service Improvement and Redesign tools. N. Improvement, Editor. 2018.
- Lohr KN. *Medicare: A Strategy for Quality Assurance*, Vol. 1. Washington, DC: National Academies Press, 1990.
- Petersen LA, Woodard LD, Urech T, Daw C, Sookanan S. Does pay-for-performance improve the quality of health care? *Ann Intern Med* 2006; **145**: 265–72.
- Roland M, Olesen F. Can pay for performance improve the quality of primary care? *BMJ* 2016; **354**: i4058.
- Ritchie C. Health care quality and multimorbidity: the jury is still out. *Med Care* 2007; **45**: 477–9.
- Institute T.J.B. Joanna Briggs Institute Reviewers' Manual, T.J.B. Institute, Editor. 2015, The University of Adelaide.

- 30 Pham MT, Rajić A, Greig JD, Sargeant JM, Papadopoulos A, McEwen SA. A scoping review of scoping reviews: advancing the approach and enhancing the consistency. *Res Synth Methods* 2014; **5**: 371–85.
- 31 Nolte E. International benchmarking of healthcare quality. A review of the literature, in Technical Reports. 2010, RAND Corporation & London School of Hygiene & Tropical Medicine: Santa Monica/London.
- 32 Ajmera M, Wilkins TL, Findley PA, Sambamoorthi U. Multimorbidity, mental illness, and quality of care: preventable hospitalizations among medicare beneficiaries. *Int J Fam Med* 2012; **2012**: 823294.
- 33 Altiner A, Schäfer I, Mellert C *et al.* Activating GENeral practitioners dialogue with patients on their Agenda (Multi-Care AGENDA) study protocol for a cluster randomized controlled trial. *BMC Fam Pract* 2012; **13**: 118.
- 34 Barbabella F, Melchiorre MG, Quattrini S, Papa R, Lamura G. *How Can eHealth Improve Care for People with Multimorbidity in Europe?* Utrecht: ICARE4EU, 2017.
- 35 Bayliss EA, McQuillan DB, Ellis JL *et al.* Using electronic health record data to measure care quality for individuals with multiple chronic medical conditions. *J Am Geriatr Soc* 2016; **64**: 1839–44.
- 36 Boyd CM, Ritchie CS, Tipton EF, Studenski SA, Wieland D. From bedside to bench: summary from the American Geriatrics Society/National Institute on Aging Research Conference on comorbidity and multiple morbidity in older adults. *Aging Clin Exp Res* 2008; **20**: 181–8.
- 37 Brilleman SL, Salisbury C. Comparing measures of multimorbidity to predict outcomes in primary care: a cross sectional study. *Fam Pract* 2013; **30**: 172–8.
- 38 Burgers JS, Voerman GE, Grol R, Faber MJ, Schneider EC. Quality and coordination of care for patients with multiple conditions: results from an international survey of patient experience. *Eval Health Prof* 2010; **33**: 343–64.
- 39 Burt J, Elmore N, Campbell SM, Rodgers S, Avery AJ, Payne RA. Developing a measure of polypharmacy appropriateness in primary care: systematic review and expert consensus study. *BMC Med* 2018; **16**: 91.
- 40 Busato A, Bhend H, Chmiel C *et al.* Improving the quality of morbidity indicators in electronic health records in Swiss primary care: a practical approach. *Swiss Med Wkly* 2012; **142**: w13611.
- 41 Bynum JPW, Chang CH, Austin A, Carmichael D, Meara E. Outcomes in older adults with multimorbidity associated with predominant provider of care specialty. *J Am Geriatr Soc* 2017; **65**: 1916–23.
- 42 Chan CL, You HJ, Huang HT, Ting HW. Using an integrated COC index and multilevel measurements to verify the care outcome of patients with multiple chronic conditions. *BMC Health Serv Res* 2012; **12**: 405.
- 43 Chandraratne NK, Pathirathna K, Harrison C, Siriwardena AN. A comparison of policies and guidelines related to multimorbidity in the UK, Australia and Sri Lanka. *Aust J Gen Pract* 2018; **47**: 14–9.
- 44 Colombo F, Garcia-Goni M, Schwierz C. Addressing multimorbidity to improve healthcare and economic sustainability. *J Comorb* 2016; **6**: 21–7.
- 45 Dantas RG, Perracini MR, Guerra RO, Ferrioli E, Dias RC, Padula RS. What are the sociodemographic and health determinants for older adults continue to participate in work? *Arch Gerontol Geriatr* 2017; **71**: 136–41.
- 46 de Bruin SR, Versnel N, Lemmens LC *et al.* Comprehensive care programs for patients with multiple chronic conditions: a systematic literature review. *Health Policy* 2012; **107**: 108–45.
- 47 Djalali S, Frei A, Tandjung R, Baltensperger A, Rosemann T. Swiss quality and outcomes framework: quality indicators for diabetes management in swiss primary care based on electronic medical records. *Gerontology* 2014; **60**: 263–73.
- 48 Drye EE, Altaf FK, Lipska KJ *et al.* Defining multiple chronic conditions for quality measurement. *Med Care* 2018; **56**: 193–201.
- 49 DuGoff EH, Dy S, Giovannetti ER, Leff B, Boyd CM. Setting standards at the forefront of delivery system reform: aligning care coordination quality measures for multiple chronic conditions. *J Healthc Qual* 2013; **35**: 58–69.
- 50 Dy SM, Pfoh ER, Salive ME, Boyd CM. Health-related quality of life and functional status quality indicators for older persons with multiple chronic conditions. *J Am Geriatr Soc* 2013; **61**: 2120–7.
- 51 Edwards ST, Landon BE. Seeking value in healthcare: the importance of generalists as primary care physicians. *J Am Geriatr Soc* 2017; **65**: 1900–1.
- 52 Eton DT, Ridgeway JL, Linzer M *et al.* Healthcare provider relational quality is associated with better self-management and less treatment burden in people with multiple chronic conditions. *Patient Prefer Adherence* 2017; **11**: 1635–46.
- 53 NQF. Multiple chronic conditions measurement framework, N.Q. Forum, Editor. 2012: Washington DC.
- 54 Fung CH, Setodji CM, Kung FY *et al.* The relationship between multimorbidity and patients' ratings of communication. *J Gen Intern Med* 2008; **23**: 788–93.
- 55 Garvey J, Connolly D, Boland F, Smith SM. OPTIMAL, an occupational therapy led self-management support programme for people with multimorbidity in primary care: a randomized controlled trial. *BMC Fam Pract* 2015; **16**: 59.
- 56 Giovannetti ER, Dy S, Leff B *et al.* Performance measurement for people with multiple chronic conditions: conceptual model. *Am J Manage Care* 2013; **19**: e359–66.
- 57 Gnädinger M, Herzig L, Ceschi A *et al.* Chronic conditions and multimorbidity in a primary care population: a study in the Swiss Sentinel Surveillance Network (Sentinella). *Int J Public Health* 2018; **63**: 1017–26.
- 58 Hadgraft NT, Willenberg L, LaMontagne AD *et al.* Reducing occupational sitting: workers' perspectives on participation in a multi-component intervention. *Int J Behav Nutr Phys Act* 2017; **14**: 73.
- 59 Haggerty JL. Ordering the chaos for patients with multimorbidity. *BMJ* 2012; **345**: e5915.
- 60 Heale R, James S, Wenghofer E, Garceau ML. Nurse practitioner's perceptions of the impact of the nurse practitioner-led clinic model on the quality of care of complex patients. *Prim Health Care Res Dev* 2018; **19**: 553–60.
- 61 Heisler M. Eliciting personal values of patients with multiple chronic conditions: why and how. *J Gen Intern Med* 2017; **32**: 1273–4.
- 62 Higashi T, Wenger NS, Adams JL *et al.* Relationship between number of medical conditions and quality of care. *N Engl J Med* 2007; **356**: 2496–504.
- 63 Hijazi HH, Alyahya MS, Hammouri HM, Alshraideh HA. Risk assessment of comorbidities on 30-day avoidable hospital readmissions among internal medicine patients. *J Eval Clin Pract* 2017; **23**: 391–401.

- 64 Holland DE, Vanderboom CE, Lohse CM *et al.* Exploring indicators of use of costly health services in community-dwelling adults with multiple chronic conditions. *Prof Case Manag* 2015; **20**: 3–11; quiz 12–3.
- 65 Huntley AL, Johnson R, Purdy S, Valderas JM, Salisbury C. Measures of multimorbidity and morbidity burden for use in primary care and community settings: a systematic review and guide. *Ann Fam Med* 2012; **10**: 134–41.
- 66 Iezzoni LI. Multiple chronic conditions and disabilities: implications for health services research and data demands. *Health Serv Res* 2010; **45**(5 Part 2): 1523–40.
- 67 Ihle A, Inauen J, Scholz U *et al.* Prospective and retrospective memory are differentially related to self-rated omission and commission errors in medication adherence in multimorbidity. *Appl Neuropsychol Adult* 2017; **24**: 505–11.
- 68 Kahn LS, Fox CH, Olawoye A, Servoss TJ, McLean-Plunkett E. Facilitating quality improvement in physician management of comorbid chronic disease in an urban minority practice. *J Natl Med Assoc* 2007; **99**: 377–83.
- 69 Kontopantelis E, Springate DA, Ashcroft DM *et al.* Associations between exemption and survival outcomes in the UK's primary care pay-for-performance programme: a retrospective cohort study. *BMJ Qual Saf* 2016; **25**: 657–70.
- 70 Krause CM, Jones CS, Joyce S *et al.* The impact of a multidisciplinary, integrated approach on improving the health and quality of care for individuals dealing with multiple chronic conditions. *Am J Orthopsychiatry* 2006; **76**: 109–14.
- 71 Leroy L, Bayliss E, Domino M *et al.* The agency for healthcare research and quality multiple chronic conditions research network: overview of research contributions and future priorities. *Med Care* 2014; **52**(3 Suppl. 2): S15–22.
- 72 Mair FS, May CR. Thinking about the burden of treatment: should it be regarded as an indicator of the quality of care? *BMJ* 2014; **349**: g6680.
- 73 Mangin D, Heath I, Jamouille M. Beyond diagnosis: rising to the multimorbidity challenge. *BMJ* 2012; **344**: e3526.
- 74 Marengoni A. Guidelines for elderly patients with multimorbidity: how to cope with a dark night without fear. *Aging Clin Exp Res* 2013; **25**: 703–5.
- 75 Mas MA. Capsule Commentary on Bennett *et al.*, engaging stakeholders to inform clinical practice guidelines that address multiple chronic conditions. *J Gen Intern Med* 2017; **32**: 918.
- 76 Melchiorre MG, Papa R, Rijken M, van Ginneken E, Hujala A, Barbabella F. eHealth in integrated care programs for people with multimorbidity in Europe: insights from the ICARE4EU project. *Health Policy* 2018; **122**: 53–63.
- 77 Min L, Kerr EA, Blaum CS, Reuben D, Cigolle C, Wenger N. Contrasting effects of geriatric versus general medical multimorbidity on quality of ambulatory care. *J Am Geriatr Soc* 2014; **62**: 1714–21.
- 78 Min LC, Wenger NS, Fung C *et al.* Multimorbidity is associated with better quality of care among vulnerable elders. *Med Care* 2007; **45**: 480–8.
- 79 Moran WP. Chaos to complexity: leveling the playing field for measuring value in primary care. *J Eval Clin Pract* 2017; **23**: 430–8.
- 80 Narayan SW, Nishtala PS. Development and validation of a medicines comorbidity index for older people. *Eur J Clin Pharmacol* 2017; **73**: 1665–72.
- 81 Ndukwe K, Burns E, Johnson J. Multiple morbidity and functional status in older adults. *J Am Geriatr Soc* 2017; **65** (Suppl 1): S248.
- 82 OECD. Patient-Reported Indicators Survey (PaRIS). 2017.
- 83 Ozminkowski RJ, Wells TS, Hawkins K, Bhattarai GR, Martel CW, Yeh CS. Big data, little data, and care coordination for medicare beneficiaries with medigap coverage. *Big Data* 2015; **3**: 114–25.
- 84 Palmer K, Marengoni A, Forjaz MJ *et al.* Multimorbidity care model: recommendations from the consensus meeting of the joint action on chronic diseases and promoting healthy ageing across the life cycle (JA-CHRODIS). *Health Policy* 2018; **122**: 4–11.
- 85 Panagioti M, Stokes J, Esmail A *et al.* Multimorbidity and patient safety incidents in primary care: a systematic review and meta-analysis. *PLoS ONE* 2015; **10**: e0135947.
- 86 Parekh AK, Goodman RA, Gordon C, Koh HK, HHS Interagency Workgroup on Multiple Chronic Conditions. Managing multiple chronic conditions: a strategic framework for improving health outcomes and quality of life. *Public Health Rep* 2011; **126**: 460–71.
- 87 Pillay M, Dennis S, Harris MF. Quality of care measures in multimorbidity. *Aust Fam Physician* 2014; **43**: 132–6.
- 88 Ricci-Cabello I, Violán C, Foguet-Boreu Q, Mounce LT, Valderas JM. Impact of multi-morbidity on quality of healthcare and its implications for health policy, research and clinical practice. A scoping review. *Eur J Gen Pract* 2015; **21**: 192–202.
- 89 Rijken M, Bekkema N, Boeckxstaens P, Schellevis FG, De Maeseneer JM, Groenewegen PP. Chronic disease management programmes: an adequate response to patients' needs? *Health Expect* 2014; **17**: 608–21.
- 90 Rijken M, Struckmann V, van der Heide I *et al.* *How to Improve Care for People with Multimorbidity in Europe?* Utrecht: ICARE4EU, 2017.
- 91 Ruscitto A, Mercer SW, Morales D, Guthrie B. Accounting for multimorbidity in pay for performance: a modelling study using UK quality and outcomes framework data. *Br J Gen Pract* 2016; **66**: e561–7.
- 92 Salisbury C. Multimorbidity: redesigning health care for people who use it. *Lancet* 2012; **380**: 7–9.
- 93 Sampalli T, Fox RA, Dickson R, Fox J. Proposed model of integrated care to improve health outcomes for individuals with multimorbidities. *Patient Prefer Adherence* 2012; **6**: 757–64.
- 94 Schiøtz ML, Høst D, Christensen MB *et al.* Quality of care for people with multimorbidity - a case series. *BMC Health Serv Res* 2017; **17**: 745.
- 95 Shadmi E, Boyd CM, Hsiao CJ, Sylvia M, Schuster AB, Boulton C. Morbidity and older persons' perceptions of the quality of their primary care. *J Am Geriatr Soc* 2006; **54**: 330–4.
- 96 Snyder L, Neubauer RL, American College of Physicians Ethics, Professionalism and Human Rights Committee. Pay-for-performance principles that promote patient-centered care: an ethics manifesto. *Ann Intern Med* 2007; **147**: 792–4.
- 97 Streit S, da Costa BR, Bauer DC *et al.* Multimorbidity and quality of preventive care in Swiss University primary care cohorts. *PLoS ONE* 2014; **9**: e96142.
- 98 Tinetti ME, Studenski SA. Comparative effectiveness research and patients with multiple chronic conditions. *N Engl J Med* 2011; **364**: 2478–81.

- 99 Uhlig K, Leff B, Kent D *et al.* A framework for crafting clinical practice guidelines that are relevant to the care and management of people with multimorbidity. *J Gen Intern Med* 2014; **29**: 670–9.
- 100 dervan HI, Snoeijs SP, Boerma WGW *et al.* *How to Strengthen Patientcentredness in Caring for People with Multimorbidity in Europe?* Utrecht: ICARE4EU, 2017.
- 101 Venkatesh A, Goodrich K, Conway PH. Opportunities for quality measurement to improve the value of care for patients with multiple chronic conditions. *Ann Intern Med* 2014; **161**(Suppl 10): S76–80.
- 102 Vetrano DL, Calderón-Larrañaga A, Marengoni A *et al.* An international perspective on chronic multimorbidity: approaching the elephant in the room. *J Gerontol A Biol Sci Med Sci* 2017; **73**: 1350–6.
- 103 Vogeli C, Shields AE, Lee TA *et al.* Multiple chronic conditions: prevalence, health consequences, and implications for quality, care management, and costs. *J Gen Intern Med* 2007; **22**(Suppl. 3): 391–5.
- 104 Wallace E, McDowell R, Bennett K, Fahey T, Smith SM. Comparison of count-based multimorbidity measures in predicting emergency admission and functional decline in older community-dwelling adults: a prospective cohort study. *BMJ Open* 2016; **6**: e013089.
- 105 Wittenberg R. The challenge of measuring multi-morbidity and its costs. *Israel J Health Policy Res* 2015; **4**: 1.
- 106 Wodchis WP. Performance measurement for people with multimorbidity and complex health needs. *Health Q* 2016; **19**: 44–8.
- 107 Zulman DM, Asch SM, Martins SB, Kerr EA, Hoffman BB, Goldstein MK. Quality of care for patients with multiple chronic conditions: the role of comorbidity interrelatedness. *J Gen Intern Med* 2014; **29**: 529–37.
- 108 Leijten FRM, Hoedemakers M, Struckmann V *et al.* Defining good health and care from the perspective of persons with multimorbidity: results from a qualitative study of focus groups in eight European countries. *BMJ Open* 2018; **8**: e021072.
- 109 Roland M, Paddison C. Better management of patients with multimorbidity. *BMJ* 2013; **346**: f2510.
- 110 Ricci-Cabello I, Stevens S, Kontopantelis E *et al.* Impact of the prevalence of concordant and discordant conditions on the quality of diabetes care in family practices in England. *Ann Fam Med* 2015; **13**: 514–22.
- 111 Akpan A, Roberts C, Bandeen-Roche K *et al.* Standard set of health outcome measures for older persons. *BMC Geriatr* 2018; **18**: 36.
- 112 Berwick DN, Cullen D, Deerberg-Wittram J *et al.* *Recommendations to OECD Ministers of Health from the High Level Reflection Group on the Future of Health Statistics. Strengthening the International Comparison of Health System Performance Through Patient-reported Indicators.* Paris: OECD, 2017.
- 113 Dowrick C. Patient-centred care for multimorbidity: an end in itself? *Lancet* 2018; **392**: 4–5.
- 114 Muth C, van den Akker M, Blom JW. The Ariadne principles: how to handle multimorbidity in primary care consultations. *BMC Med* 2014; **12**: 223.
- 115 Tinetti ME, Naik AD, Dodson JA. Moving from disease-centered to patient goals-directed care for patients with multiple chronic conditions: patient value-based care. *JAMA Cardiol* 2016; **1**: 9–10.
- 116 Starfield B. *Primary Care: Balancing Health Needs, Services and Technology.* New York: Oxford University Press, 1998.
- 117 Roland M, Nolte E. The future shape of primary care. *Br J Gen Pract* 2014; **64**: 63–4.
- 118 Nolte E. Implementing person centred approaches. *BMJ* 2017; **358**: j4126.
- 119 Coulter A. Measuring what matters to patients. *BMJ* 2017; **356**: j816.
- 120 Black N. Patient reported outcome measures could help transform healthcare. *BMJ* 2013; **346**: f167.
- 121 Valderas JM, Alonso J. Patient reported outcome measures: a model-based classification system for research and clinical practice. *Qual Life Res* 2008; **17**: 1125–35.
- 122 Edelen MO, Rose AJ, Bayliss E *et al.* Patient-reported outcome-based performance measures for older adults with multiple chronic conditions. *Rand Health Q* 2018; **8**: 3.
- 123 Ricci-Cabello I, Stevens S, Dalton ARH, Griffiths RI, Campbell JL, Valderas JM. Identifying primary care pathways from quality of care to outcomes and satisfaction using structural equation modeling. *Health Serv Res* 2018; **53**: 430–49.
- 124 Calvert M, Kyte D, Price G, Valderas JM, Hjollund NH. Maximising the impact of patient reported outcome assessment for patients and society. *BMJ* 2019; **364**: k5267.
- 125 Porter I, Evans P, Valderas JM. Routine individualized PRO feedback for the management of patients with multimorbidity in primary care: a pilot study in General Practice in England. *Qual Life Res* 2016; **25**: 1.

*Correspondence:* Jose M. Valderas, Health Services & Policy Research, University of Exeter, St Luke's Campus, Magdalen Road, Exeter EX1 2LU, UK.  
(telephone: +441392722755; e-mail: j.m.valderas@exeter.ac.uk) ■

### Supporting Information

Additional Supporting Information may be found in the online version of this article:

**Appendix.** Description of documents included in the scoping review ■